

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Potential and immuno-modulant proprieties of mesenchymal stem cells from amniotic fluid.

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1532399> since 2015-12-10T12:06:19Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Potential and immuno-modulant proprieties of mesenchymal stem cells from amniotic fluid

Katia Mareschi^{1, 2}, Deborah Rustichelli¹, Michela Muraro¹, Sara Castiglia¹, Edoardo Errichiello¹, Elena Signorino¹, Franca Fagioli¹.

¹ Stem Cell Transplantation and Cellular Therapy Unit; Pediatric Onco-Hematology Department, Regina Margherita Children's Hospital, Turin, Italy

² Department of Pediatrics of University of Turin

³ Department of Neuroscience, NIS Centre, University of Turin, Italy

⁴ Department of Gynecology and Obstetrics, University of Turin, Italy

Amniotic fluid (AF) contains stem cells which, due to their ontogenetic origin, have a high proliferative and differentiative potential and might be an attractive source of multipotent stem cells for therapeutic transplantation. We studied the phenotypic characteristics, differentiation potential and immunomodulation proprieties in vitro of mesenchymal stem cells (MSCs) isolated from human amniotic fluid (AF), during routine prenatal amniocentesis, and evaluated their compared to bone marrow (BM) MSCs.

We isolated multi-potent stem cells from AF that showed a high proliferative potential, were positive for CD90, CD105, CD29, CD44, CD73, CD166; showed Oct-4 and Nanog molecular and protein expression and differentiated into osteoblasts, adipocytes and chondrocytes. In Neural Progenitor Maintenance Medium (NPM) AF-MSCs expressed neural markers and increased Na⁺ channel density. Inactivation of the TTX-sensitive channels accelerated and became more similar to native neuronal voltage-gated Na⁺ channels.

We investigated the beneficial immunomodulatory effects of MSCs in co-cultures with different immunogenic populations such as Dendritic Cells (DCs) and regulatory CD4⁺ CD25⁺ T cells.

MSCs strongly inhibit the differentiation of monocytes to DCs and there was a significant reduction in the expression of CD83 mature DCs treated with MSCs, suggesting their skew to immature status. A decreased expression of presentation molecules (HLA-DR) and co-stimulatory molecules (CD80 and CD86) were also observed.

CD4⁺CD25⁺ regulatory T cells significantly increase in presence of MSCs. In the control group there is 5.5% of CD4⁺CD25⁺ T cells, 23.4% in co-culture with AF-derived MSCs, and 16% in co-culture with BM-derived MSCs.

These data suggest that AF-MSCs have a stronger immunosuppressive effect compared to BM-MSCs. This interesting aspect can be easily due to the more premature stemness stage of the AF-MSCs rather than BM-MSCs. We hypothesise that MSCs have proliferative, differentiative and immunological characteristics more similar to AF-MSCs than to BM-MSCs.

AF is an important multipotent stem cell source with a high proliferative potential able to originate potential precursors of functional neurons and showed a stronger immunosuppressive effect compared to BM-derived MSCs. This effect was mediated by inducing the generation of CD4⁺CD25⁺ regulatory cells and by suppressing monocyte differentiation into DCs, thus indicating the important role of AF-MSCs on immunoregulation